Preface

Complications During and After Critical Illness

Carol L. Hodgson, PhD, BAppSc(PT),
MPhil, PGDip(Cardio)
Editor

The intensive care unit (ICU) is a challenging environment for patients, families, and the staff caring for them. This is perhaps most evident when patients suffer a persistent critical illness or a prolonged stay in the ICU. These are the patients that either require prolonged mechanical ventilation or present with a cascade of new problems over time that are not necessarily related to the cause for their admission. While these patients are not common, their care is costly, increasing in prevalence, and their survival often results in long-term disability and reduced quality of life. In this issue, Drs Iwashyna and Viglianti (Michigan) discuss the epidemiology and the population-based approaches to diagnosing and treating patients who are persistently or chronically critically ill. In addition, they speculate on the future research that should be addressed in this population, including the fact that as the patient presentation changes in the ICU, there may be opportunities for improved patient care and outcomes.

With a special focus on sepsis, Dr Prescott (Michigan) highlights that one in five older patients admitted to the ICU with sepsis has a potentially preventable readmission within 90 days, most commonly for infection. In addition, she identifies the top five causes of hospital readmission following sepsis, and three common but preventable complications. She includes references, educational material, and stories for patients, families, and clinicians, including a reference to the THRIVE! (SCCM) taskforce, that seeks to improve patient and family support after critical illness by connecting with families and patients (peer support), sharing experiences and expectations of the ICU stay and the challenges after critical illness.

In addition, we investigate the new or worsening impairments in physical, psychological, and cognitive function after critical illness that persist beyond the acute hospitalization. These problems have previously been termed *post-intensive care syndrome* (PICS), a term mainly used to facilitate awareness of the functional problems faced after critical illness, prompt screening, and stimulate research. This appears to have been successful, as there were more than 200 publications using the phrase “PICS” in 2017. In this issue, we have identified and highlighted several early interventions in ICU that may have an effect on improving patient outcomes. Importantly, we have separated these to individually address neuromuscular dysfunction, psychological morbidity, or cognitive function understanding that these may occur together but often occur individually. That is, the elements of PICS do not share common signs, symptoms, or risk factors, and their management should be tailored to focus on individual pathophysiology and presentation.

Dr Brummel (Vanderbilt) highlights the heterogeneity in survivors of critical illness that limits our knowledge of outcomes. He identifies five distinct but interrelated domains that serve as a foundation for understanding functional outcomes after critical illness and describes how injuries and illnesses relate to disabilities and impairments.

Drs Montgomery, Rolfson, and Bagshaw (Alberta) reflect on the association between frailty and long-term recovery, highlighting the need to develop optimal methods for screening patients for frailty in order to identify patients who may be at increased risk of death or poor recovery. They emphasize that frail patients may need targeted interventions and specific rehabilitation, and they are likely to have increased use of health services compared with patients who are not frail.

Drs Parotto, Batt, and Herridge (Toronto) discuss the newly acquired or exacerbated organ dysfunction following critical illness that includes muscle and nerve injury that is an important determinant of long-term disability. They overview the pathophysiology and the emerging basic science of neuromuscular dysfunction and provide a rationale for future targeted clinical interventions.

Ms Paton, Dr Lane, and I (Monash University, Australia) have had the pleasure of outlining the evidence of early mobilization in ICU, including the recent concerns of implementation over evidence. There remain important questions about how to deliver early mobilization to our critically ill patients. These questions include which groups of patients are best suited to a targeted intervention of early mobilization, the optimal type of exercise prescription, and the frequency, intensity, and duration of early mobilization. We recognize that further research will be required to address these questions in future trials.

Drs Gunst and Van den Berghe (Belgium) discuss the evidence to support intensive care nutrition and recovery. They discuss the relative risks and benefits of the dose of artificial nutrition, the ideal timing of delivery of nutrition, and the controversies that surround the composition of nutrition, including immunonutrients.

Dr Girard (Pittsburgh) evaluates the evidence of delirium and sedation in the ICU setting. He emphasizes that delirium in the setting of sedation should act as a warning sign that the patient may be experiencing brain injury and the importance of eliminating sedation where possible. He acknowledges that the depth of sedation and duration of delirium are potentially modifiable and may predict long-term cognitive impairment.

Drs Gerstenblith, Davydow, and Bienvenu (Baltimore) provide an overview of the predictors of psychiatric morbidity and the experiences in ICU that may be

associated with longer-term psychiatric outcomes. They evaluate the evidence for the use of ICU diaries, while acknowledging that further work is required in this area.

Drs Haines, Quasim, and McPeake (Monash and Glasgow) discuss family and support networks in ICU, despite the lack of evidence to address how we best support patients and families. They focus on innovative approaches to build social support networks, including a model that addresses complex needs, timing of interventions, and peer support.

Carol L. Hodgson, PhD, BAppSc(PT), MPhil, PGDip(Cardio)
Australian and New Zealand
Intensive Care Research Centre
Monash University
553 St Kilda Road
Melbourne, Victoria 3004, Australia

E-mail address:

carol.hodgson@monash.edu.au